

Grape

Recommendations for Maintaining Postharvest Quality



Carlos H. Crisosto, Elizabeth J. Mitcham, and Adel A. Kader
Department of Plant Sciences, University of California, Davis

MATURITY INDICES

In California, harvest date is determined by Soluble Solids Concentration (SSC) of 14 to 17.5% depending on cultivar and production area. In some situations, the SSC/titratable acidity (TA) ratio of 20 or higher is used to determine maturity for early maturing cultivars from early production areas. For red and black colored cultivars, there is also a minimum color requirement.

QUALITY INDICES

High consumer acceptance is attained for fruit with high SSC or SSC/TA ratio. Berry firmness is also an important factor for consumer acceptance as are lack of defects such as decay, cracked berries, stem browning, shriveling, sunburned or dried berries, and insect damage.

OPTIMUM TEMPERATURE

Berry storage at -1.0 to 0°C (30-32°F) is recommended. The highest freezing point for berries is -2.7°C (27.1°F), but freezing point varies (down to -3.0°C (26.6°F)) depending on SSC. A -2.0°C (28°F) stem freezing point has been reported.

OPTIMUM RELATIVE HUMIDITY

90-95% RH and an air velocity of approximately 20-40 feet per minute (FPM) is suggested during storage.

RATES OF RESPIRATION

(Of Grape Clusters, i.e. Berries and Stems)

Temperature	ml CO ₂ /kg·hr
0°C (32°F)	1-2
5°C (41°F)	3-4
10°C (50°F)	5-8
20°C (68°F)	12-15

Stem respiration rate is approximately 15 times higher than berry respiration.

To calculate heat production, multiply ml CO₂/kg·hr by 440 to get BTU/ton/day or by 122 to get kcal/metric ton/day.


RATES OF ETHYLENE PRODUCTION

<0.1 ml/kg·hr at 20°C (68°F)

RESPONSES TO ETHYLENE

Table grapes are not very sensitive to ethylene. However, exposure to ethylene (>10 ppm) may be a secondary factor in shatter.

Produce Facts



RESPONSES TO CONTROLLED ATMOSPHERES (CA)

CA (2-5% O₂ + 1-5% CO₂) during storage/shipment is not currently recommended for table grapes because its benefit is slight and SO₂ used for decay control.

EFFECTS OF GENOTYPE ON MARKET LIFE

Market life varies among table grape cultivars grown in California and is also strongly affected by temperature management and decay susceptibility.

PHYSIOLOGICAL DISORDERS

Shatter. (Loss of berries from the cap stem) In general, shatter increases in severity with increasing maturity, i.e., the longer the fruit remains on the vine. Berries of seedless cultivars, are usually less well attached to the cap stem than seeded cultivars. Shatter varies considerably from season to season, and there is a large difference among varieties. Gibberellin applied at fruit set weakens berry attachment. Shatter occurs mainly due to rough handling during field packing with additional shatter occurring all the way to the final retail sale. Shatter incidence can be reduced by controlling pack depth and fruit packing density (cubic inches per pound), using cluster bagging, gentle handling and maintaining recommended temperature and relative humidity.

Waterberry. Waterberry is associated with fruit ripening and most often begins to develop shortly after veraison (berry softening). The earliest symptom is the development of small (1-2 mm) dark spots on the cap stems (pedicles) and/or other parts of the cluster framework. These spots become necrotic, slightly sunken, and expand to affect more areas. The affected berries become watery, soft, and flabby when ripe. In California, this disorder has been associated with a high nitrogen status vine, canopy shading, or cool weather during veraison and fruit ripening. Avoid over fertilization with nitrogen. Foliar nutrient sprays of nitrogen should be avoided in waterberry-prone vineyards. Trimming off affected berries during harvest and packing is a common practice, although labor intensive.

PATHOLOGICAL DISORDERS

Gray Mold: (*Botrytis cinerea*) Gray mold is the most destructive of the postharvest diseases of table grapes, primarily because it develops at temperatures as low as 31°F (-0.5°C) and grows from berry to berry. Gray mold first turns berries brown, then loosens the skin of the berry, its white, thread-like hyphal filaments erupt through the berry surface, and finally masses of gray colored spores develop. Wounds near harvest also provide opportunities for infections. No wound is required for infection when wet conditions occur.

Botrytis infection can be reduced by removing desiccated, infected grapes of the previous season from vines, leaf-removal canopy management, preharvest fungicides, trimming visibly infected, split, cracked, or otherwise damaged grapes before packing, prompt cooling and fumigation with sulfur dioxide (100 ppm for one hour) or use of continuous release SO₂ pads.


POSTHARVEST PHOTO GUIDE

MATURITY AND QUALITY



QUALITY

DISORDERS



STEM BROWNING



BLUE MOLD



FREEZING INJURY



GREY MOLD



SULFUR DECAY



GREY MOLD



It is the policy of the University of California not to engage in discrimination against or harassment of any person, employed by or seeking employment with the University, or in any of its programs or activities, on the basis of race, color, national origin, religion, sex, gender, gender expression, gender identity, pregnancy, physical or mental disability, medical condition (cancer-related or genetic characteristics), genetic information (including family medical history), ancestry, marital status, age, sexual orientation, citizenship, or service in the uniformed services, as well as state military and naval service. This policy is intended to be consistent with the provisions of applicable state and federal laws and University policies. University policy also prohibits retaliation against any employee or person seeking employment for bringing a complaint of discrimination or harassment pursuant to this policy. This policy also prohibits retaliation against a person who assists someone with a complaint of discrimination or harassment, or participants in any manner in an investigation or resolution of a complaint of discrimination or harassment. Retaliation includes threats, intimidation, reprisals, and/or adverse

In addition, it is the policy of the University of California to undertake affirmative action, consistent with its obligations as a Federal Contractor, for minorities and women, for persons with disabilities, and for covered veterans. The University commits itself to apply every good faith effort to achieve prompt and full utilization of minorities and women in all segments of its workforce where deficiencies exist. These efforts conform to all current legal and regulatory requirements, and are consistent with University standards of quality and excellence. In conformance with Federal regulations, written affirmative action plans shall be prepared and maintained by each campus of the University of California, by the Lawrence Berkeley National Laboratory, by the Office of the President, and by the Division of Agriculture and Natural Resources. Such plans shall be reviewed and approved by the Office of the President and the Office of the General Counsel before they are officially promulgated. Inquiries regarding the University's equal employment opportunity policies may be directed to the Affirmative Action Contact, University of California, Agriculture and Natural Resources, 2801 Second Street, Davis, CA 95618 (530) 750-1318.