

Cauliflower



Recommendations for Maintaining Postharvest Quality

Trevor V. Suslow and Marita Cantwell
Department of Plant Sciences, University of California, Davis

MATURITY INDICES

Cauliflowers are selected for size and compactness of the head or curd. Mature curds are at least 15 cm (6 inches) in diameter. Loose or protruding floral parts, creating a 'ricy' appearance, are a sign of over maturity. Cauliflower is packaged after being closely trimmed into single layer cartons of 12 to 24 heads, with 12's most common. Cauliflower is primarily marketed with closely trimmed leaves and overwrapped with perforated film. Overwraps should provide four to six 1/4-inch holes per head to allow adequate ventilation.

QUALITY INDICES

A firm and compact head of white to cream white curds surrounded by a crown of well-trimmed, turgid green leaves. Additional quality indices are size, freedom from severe yellowing due to sunlight exposure, freedom from handling defects and decay, and an absence of 'riciness'.

U.S. grade No. 1

OPTIMUM TEMPERATURE

0°C (32°F); 95-98% R.H.

Storage of cauliflower is generally not recommended for more than 3 weeks for good visual and sensory quality. Wilting, browning, yellowing of leaves, and decay are likely to increase following storage beyond 3-4 weeks or at higher than recommended storage temperatures.

RATES OF RESPIRATION


Temperature °C	Temperature °F	ml CO ₂ /kg·hr
0	32	8-9
5	41	10-11
10	50	16-18
15	59	21-25
20	68	37-42
25	77	43-48

To calculate heat production, multiply ml CO₂/kg·hr by 440 to get BTU/ton/day or by 122 to get kcal/metric ton/day.

RATES OF ETHYLENE PRODUCTION

<0.1µl /kg·hr at 20°C (68°F)

Produce Facts



RESPONSES TO ETHYLENE

Cauliflower is highly sensitive to exogenous ethylene. Discoloration of the curd and accelerated yellowing and detachment of wrapper leaf stalks will result from low levels of ethylene during distribution and short-term storage. Do not mix loads such as apples, melons and tomatoes with cauliflower.

RESPONSES TO CONTROLLED ATMOSPHERES (CA)

Controlled or modified atmospheres offer moderate to little benefit to cauliflower. Injury from low O₂ (<2%) or elevated CO₂ (>5%) may not be visual and will only be evident after cooking. When the curds become grayish, extremely soft, and emit strong off-odor. Higher levels (>10%) of CO₂ will induce this injury within 48 hours. Combined low O₂ and slightly elevated CO₂ levels (3-5%) delay leaf yellowing and the onset of curd browning by a few days.

PHYSIOLOGICAL DISORDERS

Freezing Injury. Freezing injury will be initiated at -0.8°C (30.6°F). Symptoms of freezing injury include a water soaked and greyish curd and water soaked or wilted crown leaves. The curd will become brown and gelatinous in appearance following invasion by soft-rot bacteria.

PATHOLOGICAL DISORDERS

Diseases are an important source of postharvest loss, particularly in combination with rough handling and poor temperature control. A large list of bacterial and fungal pathogens cause post-harvest losses in transit, storage, and to the consumer. Bacterial Soft-Rot (primarily *Erwinia* and *Pseudomonas*), Black Spot (*Alternaria alternata*), Grey Mold (*Botrytis cinerea*), and Cladosporium Rot are common disorders.

SPECIAL CONSIDERATIONS

For fresh-cut applications, the sensitivity of cauliflower to improper modified atmosphere (See Responses to CA) demands very careful selection of packaging films and proper temperature management.

POSTHARVEST PHOTO GUIDE

MATURITY AND QUALITY



GREEN CAULIFLOWER


DISORDERS



ETHYLENE EFFECTS



MECHANICAL DAMAGE



DISCOLORATION



It is the policy of the University of California not to engage in discrimination against or harassment of any person, employed by or seeking employment with the University, or in any of its programs or activities, on the basis of race, color, national origin, religion, sex, gender, gender expression, gender identity, pregnancy, physical or mental disability, medical condition (cancer-related or genetic characteristics), genetic information (including family medical history), ancestry, marital status, age, sexual orientation, citizenship, or service in the uniformed services, as well as state military and naval service. This policy is intended to be consistent with the provisions of applicable state and federal laws and University policies. University policy also prohibits retaliation against any employee or person seeking employment for bringing a complaint of discrimination or harassment pursuant to this policy. This policy also prohibits retaliation against a person who assists someone with a complaint of discrimination or harassment, or participants in any manner in an investigation or resolution of a complaint of discrimination or harassment. Retaliation includes threats, intimidation, reprisals, and/or adverse

In addition, it is the policy of the University of California to undertake affirmative action, consistent with its obligations as a Federal Contractor, for minorities and women, for persons with disabilities, and for covered veterans. The University commits itself to apply every good faith effort to achieve prompt and full utilization of minorities and women in all segments of its workforce where deficiencies exist. These efforts conform to all current legal and regulatory requirements, and are consistent with University standards of quality and excellence. In conformance with Federal regulations, written affirmative action plans shall be prepared and maintained by each campus of the University of California, by the Lawrence Berkeley National Laboratory, by the Office of the President, and by the Division of Agriculture and Natural Resources. Such plans shall be reviewed and approved by the Office of the President and the Office of the General Counsel before they are officially promulgated. Inquiries regarding the University's equal employment opportunity policies may be directed to the Affirmative Action Contact, University of California, Agriculture and Natural Resources, 2801 Second Street, Davis, CA 95618 (530) 750-1318.