

Chile Pepper

Recommendations for Maintaining Postharvest Quality



Marita Cantwell
Department of Plant Sciences, University of California, Davis

GENERAL INFORMATION

Chile peppers are a diverse group and come from one of 3 species of the genus *Capsicum*: *C. annuum* (most chile peppers), *C. frutescens* (Tabasco), and *C. chinense* (Habanero).

MATURITY INDICES

Mature-Green Chiles: fruit size, firmness, color
 Colored Chiles: minimum 50% coloration to achieve complete color development

QUALITY INDICES

- Uniform shape, size and color typical of variety
- Firmness
- Freedom from defects such as cracks, decay, sunburn

OPTIMUM TEMPERATURE

Chiles should be cooled as soon as possible to reduce water loss. Chile peppers are not as chilling sensitive as bell peppers. Chile peppers stored above 7.5°C (45°F) suffer more water loss, shrivel, color change and decay. Storage at 7.5°C (45°F) is considered the best for maximum shelf-life (3-5 weeks). Chiles can be stored at 5°C (41°F) for at least 2 weeks without visible signs of chilling injury. Storage at 5°C reduces water loss and shrivel, but after 2-3 weeks, chilling injury is mostly detected as discoloration of the seeds. Symptoms of chilling injury include pitting, decay, discoloration of the seed cavity, and excessive softening. Ripe or colored chiles are less chilling sensitive than mature-green chiles.

OPTIMUM RELATIVE HUMIDITY

>95%; the firmness of chile peppers is directly related to water loss during storage.


RATES OF RESPIRATION

| Temperature | 10°C (50°F) | 20°C (68°F) | 27°C (81°F) |
|---------------------------|-------------|-------------|-------------|
| ml CO ₂ /kg·hr | 5-10 | 20-30 | 40-80 |

To calculate heat production multiply ml CO₂/kg·hr by 440 to get BTU/ton/day or by 122 to get kcal/metric ton/day.

Respiration rates of chile peppers vary considerably depending on the specific type or cultivar. Jalapeno chiles have lower respiration rates while Tabasco and Chiltepin have very high respiration rates (higher than those indicated above).

Produce Facts



RATES OF ETHYLENE PRODUCTION

Some chiles such as jalapeños show a nonclimacteric physiology during color change and produce very low levels of ethylene: 0.1-0.2 $\mu\text{l/kg}\cdot\text{hr}$ at 20-25°C (68-77°F). Other chiles such as Habaneros show increases in ethylene production during ripening and may produce over 1 $\mu\text{l/kg}\cdot\text{hr}$ at 20-25°C (68-77°F).

RESPONSES TO ETHYLENE

Responses to ethylene depend on the particular variety of chile. Chile poblanos for example may respond to ethylene treatment, while Jalapeño peppers do not. As with bell peppers, holding partially colored chile peppers at warmer temperatures of 20-25°C (68-77°F) with high humidity (>95%) is effective to complete color development. Adding ethylene may further enhance ripening but response is variety dependent.

RESPONSES TO CONTROLLED ATMOSPHERES (CA)

At recommended storage temperature (7-8°C), controlled and modified atmospheres of 3-5% O₂ in combination with 0-5% CO₂ are considered to provide only slight benefit to chile peppers. Low O₂ atmospheres may retard color development. High CO₂ atmospheres (>5%) can damage mature-green chiles (pitting, discoloration, softening), while colored (fully ripe) chiles are more tolerant of CO₂.

PHYSIOLOGICAL DISORDERS

Blossom end rot. This disorder occurs as a slight discoloration or a severe dark sunken lesion at or near the blossom end; it is caused by temporary insufficiencies of calcium due to water stress and may occur under high temperature conditions when the peppers are rapidly growing.

Chilling injury. Symptoms of chilling injury include surface pitting, water-soaked areas, decay (especially *Alternaria*), and discoloration of the seed cavity.

PATHOLOGICAL DISORDERS

Common decay causing organisms are *Botrytis*, *Alternaria*, and soft rots of fungal and bacterial origin.

Botrytis or Grey mold decay is a common decay-causing organism on chile peppers; field sanitation and prevention of wounds on the fruit help reduce its incidence. *Botrytis* will grow well at the recommended storage temperatures. Hot water dips of peppers can effectively control botrytis rot (55°C [130°F] water for 4 minutes) without causing fruit injury.

Bacterial Soft Rot. Soft rotting areas can be caused by several bacteria which attack damaged tissue; soft rots can also be common on washed or hydrocooled chile peppers where water sanitation was deficient.

Mechanical damage (crushing, stem punctures, scrapes, etc.) is common on chile peppers; physical injury not only detracts from the visual quality of the chiles but also causes increased

SPECIAL CONSIDERATIONS

The pungency or hotness of chile peppers is due to capsaicinoids (capsaicin is the main one) and hotness varies depending on the cultivar or genetics of the chile peppers. Environmental factors and maturity of the chiles also affects the capsaicin concentrations. Chiles that are stored under a range of temperatures and are still of marketable quality maintain their capsaicin concentrations.

For U.S. markets, corking (corky striations on the fruit surface) is considered unattractive. However in other markets, corking is a recognized characteristic of certain cultivars and is usually associated with jalapeño chiles that are prepared in oil and vinegar as a side dish.

POSTHARVEST PHOTO GUIDE

DISORDERS



Botrytis decay on stored Jalapeño chiles




GREY MOLD



BLOSSOM END ROT



CHILLING INJURY



It is the policy of the University of California not to engage in discrimination against or harassment of any person, employed by or seeking employment with the University, or in any of its programs or activities, on the basis of race, color, national origin, religion, sex, gender, gender expression, gender identity, pregnancy, physical or mental disability, medical condition (cancer-related or genetic characteristics), genetic information (including family medical history), ancestry, marital status, age, sexual orientation, citizenship, or service in the uniformed services, as well as state military and naval service. This policy is intended to be consistent with the provisions of applicable state and federal laws and University policies. University policy also prohibits retaliation against any employee or person seeking employment for bringing a complaint of discrimination or harassment pursuant to this policy. This policy also prohibits retaliation against a person who assists someone with a complaint of discrimination or harassment, or participants in any manner in an investigation or resolution of a complaint of discrimination or harassment. Retaliation includes threats, intimidation, reprisals, and/or adverse

In addition, it is the policy of the University of California to undertake affirmative action, consistent with its obligations as a Federal Contractor, for minorities and women, for persons with disabilities, and for covered veterans. The University commits itself to apply every good faith effort to achieve prompt and full utilization of minorities and women in all segments of its workforce where deficiencies exist. These efforts conform to all current legal and regulatory requirements, and are consistent with University standards of quality and excellence. In conformance with Federal regulations, written affirmative action plans shall be prepared and maintained by each campus of the University of California, by the Lawrence Berkeley National Laboratory, by the Office of the President, and by the Division of Agriculture and Natural Resources. Such plans shall be reviewed and approved by the Office of the President and the Office of the General Counsel before they are officially promulgated. Inquiries regarding the University's equal employment opportunity policies may be directed to the Affirmative Action Contact, University of California, Agriculture and Natural Resources, 2801 Second Street, Davis, CA 95618 (530) 750-1318.